

RF2112 / 061 / PCT

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
15 February 2001 (15.02.2001)

PCT

(10) International Publication Number
WO 01/10814 A1

- (51) International Patent Classification⁷: C07C 203/04 (74) Agents: SAMA, Daniele et al.; Sama Patents, Via G.B. Morgagni, 2, I-20129 Milano (IT).
- (21) International Application Number: PCT/EP00/07222
- (22) International Filing Date: 27 July 2000 (27.07.2000) (81) Designated States (national): AE, AL, AU, BA, BB, BG, BR, CA, CN, CR, CU, CZ, DM, EE, GD, GE, HR, HU, ID, IL, IN, IS, JP, KP, KR, LC, LK, LR, LT, LV, MA, MG, MK, MN, MX, NO, NZ, PL, RO, SG, SI, SK, TR, TT, UA, US, UZ, VN, YU, ZA.
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data: MI99A001753 4 August 1999 (04.08.1999) IT (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- (71) Applicant (for all designated States except US): NICOX S.A. [FR/FR]; 45, avenue Kléber, F-75116 Paris (FR).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): BENEDINI, Francesca [IT/IT]; Via Padova, 286, I-20100 Milano (IT). OLDANI, Erminio [IT/IT]; Via San Massimo, 82, I-20018 Sedriano (IT). CASTALDI, Graziano [IT/IT]; Via Livia Gallina, 5, I-28072 Briona (IT). TARQUINI, Antonio [IT/IT]; Via Postumia, 23/2, I-15057 Tortona (IT).

Published:

— With international search report.

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

WO 01/10814 A1

(54) Title: PROCESS FOR THE PREPARATION OF NAPROXENE NITROXYALKYLESTERS

(57) Abstract: A process for obtaining nitroxyalkylesters of the 2-(S)-(6-methoxy-2-naphthyl)-propanoic acid having an enantiomeric excess higher than or equal to 95 %, preferably higher than or equal to 98 %, characterized in that an halide of the 2-(S)-(6-methoxy-2-naphthyl)-propanoic acid of formula A-Hal, wherein A is the acid acyl residue, is reacted in an inert organic solvent with an aliphatic nitroxyalkanol HO-Y-ONO₂, wherein Y is a C₂-C₂₀ alkylene or a cycloalkylene from 3 to 8 carbon atoms, or an alkylene as defined containing a cycloalkylene as defined, in the presence of an inorganic base.